

Table VI. Comparison of nutritional biochemistry (fasting) of Spata and Melbourne samples (mean \pm SD)

	Spata		Melbourne	
	70-79 years	80+ years	70-79 years	80+ years
<i>Men</i> (n)	24	14	47	14
Serum albumin (g/l)	44.5 \pm 2	45.0 \pm 3	43.8 \pm 3 ^c	43.9 \pm 2 ^d
Total lymphocyte count (mm ³)	2180 \pm 610 ⁱ	2227 \pm 429	1813 \pm 548 ^{ci}	1914 \pm 395
% lymphocytes (tlc/wbc \times 100)	33 \pm 12 ^a	35 \pm 5 ^l	29 \pm 9 ^c	27 \pm 5 ^j
Haemoglobin (g/dl)	14.2 \pm 1 ^{ai}	14.3 \pm 2 ^b	14.9 \pm 1 ^{ci}	14.5 \pm 1
Haematocrit (%)	44.4 \pm 4 ^a	44.5 \pm 4 ^b	45.1 \pm 6 ^c	41.1 \pm 9
Plasma iron (μ mol/l)	23.0 \pm 7	24.9 \pm 6	21.7 \pm 7	20.6 \pm 7 ^d
Total iron binding (μ mol/l)	59.9 \pm 10	57.9 \pm 7	62.4 \pm 11	62.5 \pm 11
Transferrin saturation (%)	39.8 \pm 16	43.9 \pm 12 ^b	35.5 \pm 14 ^c	35.3 \pm 12 ^d
Plasma ferritin (μ g/l)	96.9 \pm 88 ⁱ	76.7 \pm 39 ^j	184.4 \pm 126 ⁱ	167 \pm 139 ^j
Plasma folate (nmol/l)	19.8 \pm 11	17.3 \pm 5	17.3 \pm 7	18.4 \pm 7
Plasma B ₁₂ (pmol/l)	284.9 \pm 153	244.4 \pm 141 ^b	286.6 \pm 221 ^f	239 \pm 102 ^f
Serum cholesterol (mmol/l)	6.1 \pm 1 ^a	6.6 \pm 2	6.1 \pm 1	6.1 \pm 1
Serum triglycerides (mmol/l)	1.6 \pm 0.7	1.7 \pm 1	1.3 \pm 0.7	1.3 \pm 0.5
Serum HDL-cholesterol (mmol/l)	1.4 \pm 0.3 ^{ai}	1.3 \pm 0.2 ^b	1.2 \pm 0.3 ^{ci}	1.4 \pm 0.3
Serum LDL-cholesterol (mmol/l)	3.9 \pm 1	4.5 \pm 1	4.2 \pm 1	4.1 \pm 1
LDL/HDL	2.9 \pm 1 ⁱ	3.5 \pm 1	3.6 \pm 1 ^{ci}	3.1 \pm 1
Fasting plasma glucose (mmol/l)	5.5 \pm 1	5.5 \pm 0.7	6.3 \pm 2	6.0 \pm 2
<i>Women</i> (n)	17	8	32	15
Serum albumin (g/l)	44.3 \pm 2 ^k	46.7 \pm 3 ^{gl}	42.3 \pm 2.5 ^{ck}	41.3 \pm 2 ^{dl}
Total lymphocyte count (mm ³)	2198 \pm 447	2036 \pm 507	2162 ^c \pm 550	1928 \pm 526
% lymphocytes (tlc/wbc \times 100)	36 \pm 4 ^{ak}	35 \pm 12	33 \pm 6 ^{chk}	27 \pm 7 ^h
Haemoglobin (g/dl)	12.7 \pm 0.6 ^{ak}	13.0 \pm 0.6 ^b	13.5 \pm 1 ^{ck}	13.7 \pm 2
Haematocrit (%)	39.9 \pm 2 ^a	41.0 \pm 2 ^b	40.3 \pm 3 ^c	40.9 \pm 4
Plasma iron (μ mol/l)	20.8 \pm 6	20.9 \pm 6 ^l	18.8 \pm 6	15.9 \pm 5 ^{di}
Total iron binding (μ mol/l)	62.2 \pm 10	62.4 \pm 8	64.8 \pm 9.5	62.2 \pm 8
Transferrin saturation (%)	34.8 \pm 13	34.1 \pm 10 ^{bl}	29.8 \pm 10 ^c	25.8 \pm 8 ^{di}
Plasma ferritin (μ g/l)	72.3 \pm 75	66.5 \pm 35	129.5 \pm 122	102.0 \pm 71
Plasma folate (nmol/l)	21.8 \pm 11	21.9 \pm 7 ^l	19.2 \pm 8.1	15.9 \pm 4.7 ^l
Plasma B ₁₂ (pmol/l)	439.5 \pm 390	580.4 \pm 541 ^{bl}	310.6 \pm 176	252.7 \pm 93 ^l
Serum cholesterol (mmol/l)	7.1 \pm 2.5 ^a	7.2 \pm 1.3 ^l	6.3 \pm 1.2	6.0 \pm 1.3
Serum triglycerides (mmol/l)	1.3 \pm 0.7	1.6 \pm 0.5	1.3 \pm 0.5	1.4 \pm 0.6
Serum HDL-cholesterol (mmol/l)	1.7 \pm 0.4 ^a	1.5 \pm 0.3 ^b	1.5 \pm 0.4 ^c	1.3 \pm 0.2
Serum LDL-cholesterol (mmol/l)	4.8 \pm 2.4	5.0 \pm 1.3	4.2 \pm 1.0	4.1 \pm 1.3
LDL/HDL	2.9 \pm 1.5	3.5 \pm 1.5	2.9 \pm 1.0 ^c	3.2 \pm 1.2
Fasting plasma glucose (mmol/l)	5.8 \pm 1.7	5.8 \pm 1.3	5.9 \pm 2.4	6.7 \pm 2.7

Pairs of letters indicate significant differences, Wilcoxon $p < 0.05$:

a,b,c or d within centres—between sexes for a given age group;

e,f,g or h within centres—between age groups for a given sex;

i,j,k or l between centres—for a given age group and sex.

reported a similarly high prevalence of heart trouble (23%) comparable with the prevalence reported in the elderly Anglo-Celtic Australian study [22] and elderly Greek subjects in the Euronut-Seneca study [15]. In contrast with the men, Greek women in Melbourne (mainly aged 80+) had a significantly greater prevalence of heart disease (40%) than the men, the Spata women (16%) and the women in the elderly Anglo-Celtic Australian study [22]. Melbourne and Spata men also reported a similarly low prevalence of cancer (4%, excluding skin cancer) which was significantly lower than the prevalence reported in the elderly Anglo-Celtic Australian study (>7%, excluding skin cancer) but comparable with the prevalence reported in elderly

Greeks in the Euronut study [15]. Melbourne women (mainly aged 80+) had a greater prevalence of cancer than the men and the Spata women and the women in the Anglo-Celtic Australian study. This suggests that Melbourne women are losing their protection against heart disease, and also cancer, at a faster rate than the men and the Spata women.

The traditional Greek diet can be defined as vegetarian, foods from plant sources forming the core of the diet, while foods from animal sources provide the fringe of the diet (plant to animal food ratio 75 : 25) [17, 48, 49]. This dietary pattern, prevalent prior to the 1960s, has been associated with a lower prevalence of heart disease, colonic cancer [5, 6, 42] and reduced risk

Table VII. Comparison of anthropometric measurements of Spata and Melbourne samples (mean \pm SD)

	Spata		Melbourne	
	70-79 years	80+ years	70-79 years	80+ years
<i>Men</i> (n)	26	15	64	28
Height (cm)	165.9 \pm 6 ^a	163.9 \pm 6 ^b	165.2 \pm 6 ^c	163.2 \pm 6 ^d
Weight (kg)	75.7 \pm 14 ^a	67.9 \pm 10	76.3 \pm 11 ^c	72.3 \pm 11 ^d
BMI (kg/height m ²)	27.4 \pm 4	25.2 \pm 3	27.9 \pm 4 ^c	27.0 \pm 4
% <20	3.8	6.6	1.6	3.5
% 20-24.9	26.9	26.7	20.3	28.6
% 25-29.9	34.7	66.7	48.4	39.3
% \geq 30	34.6 ^{ac}	0.0 ^{bej}	29.7 ^c	28.6 ^{dj}
Estimated body fat (%)*	32 \pm 5 ^a	32 \pm 4 ^b	33 \pm 4 ^c	34 \pm 4 ^d
Estimated lean mass (kg)*	51 \pm 6 ^{ac}	46 \pm 5 ^{bc}	51 \pm 5 ^{cf}	47 \pm 6 ^{df}
Arm muscle area (cm ²)	51.8 \pm 9 ^{ac}	45.2 \pm 6.6 ^c	56.2 \pm 12 ^{cf}	52.0 \pm 9 ^{df}
Umbilical circumference	100.8 \pm 11	96.9 \pm 8.9 ^b	100.1 \pm 9 ^c	100.5 \pm 10
Maximal gluteal circumference	105.5 \pm 9	102.7 \pm 8.1	103.0 \pm 7	101.1 \pm 7
Umbilical/maximal gluteal	0.95 \pm 0.04 ^a	0.94 \pm 0.04 ^{bj}	0.98 \pm 0.06 ^c	0.99 \pm 0.06 ^j
% \geq 0.9	96.2	86.7	93.7	100.0
<i>Women</i> (n)	20	9	59	35
Height (cm)	151.3 \pm 6 ^a	149.8 \pm 4 ^b	149.8 \pm 5 ^c	148.5 \pm 6 ^d
Weight (kg)	64.7 \pm 10 ^a	61.2 \pm 8	68.8 \pm 11 ^{ch}	61.4 \pm 14 ^{dh}
BMI	28.2 \pm 4	27.2 \pm 4	30.7 \pm 5 ^{ch}	27.8 \pm 6 ^h
% <20	0.0	0.0	1.7	8.5
% 20-24.9	15.0	44.5	3.4	28.6
% 25-29.9	60.0	22.2	45.8	22.9
% \geq 30	25.0 ^{agk}	33.3 ^{agl}	49.1 ^{chk}	40.0 ^{dhl}
Estimated body fat (%)*	47 \pm 3 ^a	49 \pm 2 ^b	48 \pm 3 ^c	50 \pm 3 ^d
Estimated lean mass (kg)*	34 \pm 5 ^a	31 \pm 4 ^b	35 \pm 5 ^{ch}	30 \pm 6 ^{dh}
Arm muscle area (cm ²)	2.8 \pm 10 ^{agk}	40.3 \pm 6 ^g	49.7 \pm 11 ^{chk}	42.0 \pm 12 ^{dh}
Umbilical circumference	107.0 \pm 11	105.7 \pm 12 ^b	108.0 \pm 10 ^c	103.6 \pm 13
Maximal gluteal circumference	107.3 \pm 10	104.9 \pm 9	105.6 \pm 9	102.1 \pm 9
Umbilical/maximal gluteal	0.99 \pm 0.03 ^a	1.01 \pm 0.08 ^b	1.02 \pm 0.06 ^c	0.98 \pm 0.17
% \geq 0.8	100.0	100.0	100.0	100.0

Pairs of letters indicate significant differences, χ^2 (%) or Wilcoxon (score), $p < 0.05$:

a,b,c or d within centres—between sexes for a given age group;

e,f,g or h within centres—between age groups for a given sex;

i,j,k or l between centres—for a given age group and sex.

* Duerenberg equation (see Methods).

of death in later life [12], and is now regarded as a prudent diet [17, 49, 50]. The traditional Greek diet translates into the following macronutrient proportions; low-moderate protein intake (<15% of energy intake); high total fat intake (40-45%); high mono-unsaturated fat intake (20-22%); low saturated fat intake (10-12%); low polyunsaturated fat intake (<5%); moderate carbohydrate intake (38-44%; complex 28%; refined 12%) and low-moderate alcohol intake (<5%) [16, 17]. In recent years, Greeks in Greece [21] and abroad [2, 4, 5] have been moving away from the traditional dietary pattern, towards a more 'affluent' diet, characterized by a proportionately higher animal food intake (especially marked in migrants) and lower plant food intake. Since Spata was chosen for its 'traditionality' it was not surprising that the plant to animal food ratio was significantly greater in Spata (69:31) than in Melbourne (65:35), with the latter approaching the lower ratio found

amongst elderly Anglo-Celtic Australians (59:41) [22]. Energy intakes from the macronutrients were in the realm of the traditional Greek diet for both locations, except that Melbourne subjects had lower intakes of complex carbohydrates (37%) and greater intakes of refined carbohydrates (14%; mainly from fruit juice), protein (19%; mainly from meat) and polyunsaturated fat (6%; mainly from margarine), similar to levels found in Anglo-Celtic Australians [22, 49]. The greater intakes of meat (130 g) and protein (19% energy) by Melbourne Greeks are approaching the intakes of elderly Anglo-Celtic Australians (160 g/day; 19% energy) [22]. Similar findings have been reported in other studies [4, 6]. Even though meat intake was significantly lower in Spata (90 g/day), this was still greater than intakes reported in Greece in the 1960s (35 g/day) [44]. In contrast, fish consumption has remained high in both Melbourne and Spata (50 g/day) since the 1960s (40 g/day) and has not dropped to the

lower levels reported in Anglo-Celtic Australians (15 g/day) [22, 51]. National authorities are now inclined to recommend decreased consumption of animal protein in preference for plant proteins, which have been associated with reduced rates of heart disease and colonic cancer [52, 53]. Continued high intake of animal proteins in Melbourne Greeks may be contributing to the changing prevalence of these diseases. Melbourne Greek elderly subjects (especially men) have also maintained the high vegetable (400 g/day) and legume (70 g/day) intake of the 1960s [51] but have markedly decreased their consumption of cereals (250 g/day), complex carbohydrates (22% energy intake), and fruit (200 g/day). Elderly Anglo-Celtic Australians have high plant food intakes similar to Melbourne Greeks, but the types of plant foods consumed are significantly different—namely a lower intake of legumes (20 g/day, comprised mainly of peas) and cereals (comprised mainly of bread and breakfast cereals). Certain legumes and cereals have been shown to lower cholesterol and to be potentially protective against colonic cancer [53, 54]. Melbourne Greeks may have continued protection against colon cancer relative to Anglo-Celtic Australians owing to their higher intakes of specific legumes (e.g. dried beans, chick-peas) and cereals (rice, pasta).

The high intake of total fat by elderly Greeks (40–42% energy intake) is also in agreement with other studies in Greece [15–17]. Case-control studies from Greece have not shown adverse effects of the currently high fat intake (40–45%) on risk of cancer, coronary heart disease or obesity. Researchers in Greece argue that the current evidence does not justify reduction of fat intake in the Greek diet if derived mainly from mono-unsaturated fats [16, 41, 43], particularly since olive oil facilitates increased consumption of plant foods [12] and has been associated with reduced rates of heart disease [54]. Olive oil consumption has been reported to have decreased markedly in Greece since the 1960s (from about 60 g/day [51, 55] and to have decreased even further in migrant Greeks [5, 6]. Much of this olive oil has been replaced by margarines and other polyunsaturated oils, which have recently been linked to heart disease and cancer [53]. Spata elderly subjects consumed only olive oil (30 g/day) which was an important marker of adherence to the traditional Greek diet. In the Euronut study, Greek elderly subjects in a rural village reported eating about 30 g/day [12]. In contrast, Melbourne elderly consumed significantly less olive oil (18 g/day) and had introduced margarine into their diets (3 g/day), but not to the levels found in elderly Anglo-Celtic Australians (15 g/day) [22]. Red wine has also been associated with reduced incidence and mortality from coronary heart disease [56], whereas beer has been epidemiologically linked with rectal cancer [57] and heart disease [50, 58]. About 90% of the alcohol consumed by Spata men was from locally produced wine (170 g/day; 60% red, 40% white). In Melbourne, only 58% of the alcohol consumed by men was from wine (70 g/day); 37% was as beer (50 g/day) and 5% as spirits. The reduced intake of red wine

and olive oil and the increased intake of margarine and beer may be playing a significant role in the changing health profile of migrant Greeks. With respect to micronutrient intake, the diets of Melbourne Greeks were more nutritionally adequate than Spata Greeks, owing to a significantly greater intake of meat, vegetables, legumes and milk. The nutrients at greatest risk of deficiency included thiamin, magnesium, and vitamin A, with more than 50% of Spata elderly subjects failing to achieve two-thirds of the RDI compared with less than 20% of Melbourne subjects. The risk of deficiency was lower for calcium, followed by zinc and lastly riboflavin, in both study sites. Vitamin C, niacin, iron, potassium (except for Spata women) and phosphorus were least likely to be inadequately consumed, especially in Melbourne. Published studies of elderly people have also reported similarly low intakes of these nutrients [15, 59, 60]. Measurements of body fatness and nutritional biochemistry revealed some important differences between the two locations. Melbourne Greeks (particularly the women) were at greater risk of disease than Spata Greeks because of the greater prevalence of morbid obesity, centrally distributed fat [61], unfavourable blood lipid profiles and impaired immunity [62]. The immune function of Melbourne Greeks appears to be dropping to the lower levels found in Anglo-Celtic Australians [22]. High levels of storage iron have been associated with impaired immunity, as well as heart disease and colon cancer [63]. High storage iron levels were more prevalent in Melbourne (20%) than Spata (3%), approaching the high levels found in Anglo-Celtic Australians (>20%). The extent to which iron status contributes to the deteriorating health and immune function of elderly Greeks needs further investigation.

In conclusion, there is evidence that elderly Spata Greeks (mainly men aged 70–79) are 'healthier' than elderly Melbourne Greeks, especially with respect to blood lipid profiles and immune function. Compared with Spata Greeks, Melbourne Greeks had significantly greater intakes of animal foods (meat), legumes, protein, margarine, polyunsaturated fats, and beer, and lower intakes of cereals, carbohydrates, wine and olive oil. The contribution of these dietary differences, as well as the influence of high storage iron levels, impaired immunity and greater prevalence of obesity and abdominal fatness, to the increasing prevalence of heart disease and cancer, especially amongst women, merits further study.

Acknowledgements

This study was funded in part by the Sandoz Foundation for Gerontological Research. The statistical assistance of Dr Bridget Hsu-Hage and the co-operation of the staff at the Spata Community Health Centre are gratefully acknowledged.

References

1. Young C. *Selection and survival: immigrant mortality in Australia*. Studies in Adult Migrant Education. Depart-

- ment of Immigration and Ethnic Affairs. Australian Government Printing Services, Canberra, Australia, 1986.
2. Powles JW. Best of both worlds? Attempting to explain the persisting low mortality of Greek migrants to Australia. In: Caldwell JC, ed. *Health transition: cultural, social and behavioural determinants of health: what is the evidence?* Proceedings of a workshop, Canberra 15–19 May 1989. Australian National University, Canberra, 1990.
 3. World Health Organization. *Health statistics annual 1991*. Geneva, 1992.
 4. Powles JW, Hage BH, Cosgrove M. Health related expenditure patterns in selected migrant groups: data from the Australian Household Expenditure Survey, 1984. *Community Health Stud* 1990;14:1–7.
 5. Powles JW, Hage BH, Ktenas DK. Who came? Who stayed behind? Selection and migration from the Greek island of Levkada. *Departmental research report no. 88/1*. Department of Social and Preventive Medicine, Monash University, Melbourne, Australia, 1988.
 6. Powles JW, Ktenas D, Sutherland C, Hage B. Food habits in Southern-European migrants: a case study of migrants from the Greek island of Levkada. In: Truswell S, Wahlqvist M, eds. *Food habits in Australia*. Balwyn, Australia: Rene Gordon, 1988;201–23.
 7. McMichael AJ, McCall MG, Hartshorne JM, Woodings TL. Patterns of gastro-intestinal cancer in European migrants to Australia: role of dietary change. *Int J Cancer* 1980;25:431.
 8. Australian Bureau of Statistics. *National Health Survey, summary of results 1989–90*. ABS cat. no. 4364.0, Australian Government Printing Services, Canberra, Australia, 1991.
 9. Australian Council on the Ageing and Department of Community Services. *Older people at home, a report of a 1981 joint survey conducted in Melbourne and Adelaide*. Australian Government Printing Services, Canberra, Australia, 1986.
 10. Australian Institute of Multicultural Affairs. *Community and institutional care for aged migrants in Australia: research findings*. AIMA, Melbourne, Australia, 1986.
 11. National Heart Foundation. *Risk factor prevalence study no. 2, 1983*. National Heart Foundation of Australia, Canberra, Australia, 1985.
 12. Trichopoulou A, Kouris-Blazos A, Vassilakou T, et al. The diet and survival of elderly Greeks: a link to the past. *Am J Clin Nutr* 1995;61(6S):1346–50.
 13. Kromhout D, Keys A, Aravanis C, et al. Food consumption patterns in the 1960s in seven countries. *Am J Clin Nutr* 1989;49:889–94.
 14. Kafatos A, Kouroumalis I, Vlachonikolis I, Theodorou C, Labadarios D. Coronary heart disease risk-factor status for the Cretan urban population in the 1980s. *Am J Clin Nutr* 1991;54:591–8.
 15. de Groot LCPGM, van Staveren WA, Hautvast JGAJ. Euronut-Seneca, nutrition and the elderly in Europe: a concerted action on nutrition and health in the European Community. *Eur J Clin Nutr* 1991;45(suppl 3):5–185.
 16. Trichopoulou A, Toupadaki N, Tzonou A, et al. The macronutrient composition of the Greek diet: estimates derived from six case-control studies. *Eur J Clin Nutr* 1993;47:549–58.
 17. Trichopoulou A, Katsouyanni K, Gnardellis Ch. The traditional Greek diet. *Eur J Clin Nutr* 1993;47(suppl 1):76–81.
 18. Rutishauser I, Wahlqvist ML. Food intake patterns of Greek migrants to Melbourne in relation to duration of stay. *Proc Nutr Soc Aust* 1983;49–55.
 19. Kosmidis G, Rutishauser I, Wahlqvist ML, McMichael A. Food intake patterns amongst Greek migrants in Melbourne. *Proc Nutr Soc Aust* 1980;5:165.
 20. Department of Community Services and Health. *National Dietary Survey of Adults, 1983*. Canberra, Australian Government Publishing Service, 1987.
 21. Trichopoulou A, Efstathiadis PP. Changes in nutrition patterns and health indicators at the population level in Greece. *Am J Clin Nutr* 1989;49:1042–7.
 22. Wahlqvist ML, Davies L, Hsu-Hage BH-H, et al. *Cross-cultural study of food habits and health status in later life: description of elderly communities and lessons learnt*. *Asia Pacific J Clin Nutr* and United Nations University Press, 1995.
 23. Wahlqvist ML, Kouris A. Trans-cultural aspects of nutrition in old age. *Age Ageing* 1990;19(suppl):S43–52.
 24. Wahlqvist ML, Hsu-Hage B, Kouris-Blazos A, Lukito W. Food habits in later life: a cross-cultural study. In: Wahlqvist M, et al. eds. *Nutrition in a sustainable environment*. Proceedings of the XVth International Congress of Nutrition, United Kingdom: Smith-Gordon, 1994;345–53.
 25. Wahlqvist ML, Kouris A, Davies L, Scrimshaw N. Development of a Survey Instrument for the Assessment of Food Habits and Health in Later Life. In: Moyal M, ed. *Dietetics in the 90s: role of the dietitian/nutritionist*. Proceedings of the International Congress of Dietetics, Paris: J Libbey Eurotext, 1988.
 26. Kouris A, Wahlqvist ML, Trichopoulou A, Polychronopoulos E. Food habits and health of elderly in Spata, Greece: application of Survey Instrument. In: *New global era: global harmony through nutrition*. Proceedings of the XIVth International Congress of Nutrition, United Kingdom: Smith-Gordon, 1989.
 27. Kouris-Blazos A. Elderly Greeks in Spata, Greece and Melbourne, Australia: food habits, health and lifestyle. PhD thesis. Monash University, Melbourne, Australia, 1994.
 28. Kouris A, Wahlqvist ML, Trichopoulou A, Polychronopoulos E. Use of combined methodologies in assessing food beliefs and habits of elderly Greeks in Greece. *Food Nutr Bull* 1991;13:139–44.
 29. Australian Bureau of Statistics. *The overseas-born in Victoria—a statistical profile*, 1986. ABS cat. no. 2503.2, Australian Government Printing Services, Canberra, Australia, 1991.
 30. Lawton MP, Moss M, Fulcomer M, Kleban MH. A research and service oriented multilevel assessment instrument. *J Gerontol* 1982;37:91–9.
 31. Fillenbaum, GG. *The wellbeing of the elderly: approaches to multidimensional assessment*. Geneva: World Health Organization, Offset, Publication no. 84, 1984.
 32. Macrae FA, Lambert JR, Wahlqvist MC, Selbie L, Brouwer R, MacLennan R. The Australian Polyp Prevention Project: compliance to randomized dietary intervention. *Proc Gastroenterol Soc Aust* 1989.
 33. Trichopoulou A, Kampman E, Koliass E, Georga K. *A photographic method to estimate food and nutrient intake in Greece—user's manual*. Department of Nutrition and Biochemistry, Athens School of Public Health, Greece, 1988.
 34. Schofield WN, Schofield C, James WPT. Basal metabolic

- rate-review, prediction, together with annotated bibliography source material. *Hum Nutr Clin Nutr* 1985;39C (suppl 1):1-96.
35. Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Prentice AM. Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* 1991;45:569.
 36. FAO/WHO/UNU. *Energy and protein requirements*. Report of a joint FAO/WHO/UNU Expert consultation. WHO Tech Rep Ser No. 74 Geneva: World Health Organization, 1985.
 37. National Research Council. *Recommended dietary allowances*. 10th edn. Washington DC: National Academy Press, 1989.
 38. Deurenberg P, van der Kooy K, Leenen R, Weststrate JA, Seidell JC. Sex and age specific prediction formulas for estimating body composition from bioelectrical impedance: a cross-validation study. *Int J Obesity* 1991;15:17-25.
 39. Gibson R. *Principles of nutritional assessment*. Oxford: Oxford University Press, 1990.
 40. Keys A, ed. *Seven countries: a multivariate analysis of death and coronary heart diseases*. Cambridge, Mass.: Harvard University Press, 1980.
 41. Trichopoulou A. Mediterranean diet, disease and nutrition guidelines. In: Benito E, Giacosa A, Hill MJ, eds. *Public education on diet and cancer*. (Proc 9th Annu Symp Eur Organ Coop Cancer Prev Stud) Kluwer Academic, 1992;135-42.
 42. Kafatos A, Kouroumalis I, Vlachonikolis I, Theodorou C, Labadarios D. Coronary heart disease risk-factor status for the Cretan urban population in the 1980s. *Am J Clin Nutr* 1991;54:591-8.
 43. Trichopoulos D, Skalkidis Y, Trichopoulou A, Touloumi O, Souliou L. The health of the Greek population: yesterday, today, tomorrow. *Mater Med Greca* 1989;315.
 44. Aravanis C, Ioannidis PJ. Nutritional factors and cardiovascular diseases in the Greek Islands Heart Study. In: Lovenberg W, Yamori Y, eds. *Nutritional prevention of cardiovascular disease*. New York: Academic Press, 1984;125-35.
 45. Trichopoulos D, Xirouchaki E, Tzonou A, Katsouyanni K, Zavitsanos X. Cancer in Greece: regional differences and secular trends. *Mater Med Greca* 1982;10:20-30.
 46. Spiller GA. *The Mediterranean diets in health and disease*. Health Research and Studies Centre, Los Altos, California and Sphera Foundation, New York: Van Nostrand Reinhold, 1991.
 47. Wahlqvist ML, Kouris-Blazos A, Trichopoulou A, Polychronopoulos E. The wisdom of the Greek cuisine and way of life: comparison of the food and health beliefs of elderly Greeks in Greece and Australia. *Age Nutr* 1991;2:163-73.
 48. Harvard School of Public Health. *International conference on the diets of the Mediterranean*. Cambridge, Mass.: Oldways Preservation and Exchange Trust Publications, 1993.
 49. James WPT, Duthie G, Wahle K. The Mediterranean diet: protective or simply non-toxic? *Eur J Clin Nutr* 1989;43(supp 2):31-41.
 50. Baghurst KI, Crawford DA, Worsley A, Syrette JA, Record SJ, Baghurst PA. The Victorian Nutrition Survey: a profile of the energy, macronutrient and sodium intakes of the population. *Community Health Stud* 1988;22:42-54.
 51. Food and Agricultural Organization. *Standardized food balance sheets*. Rome: FAO, 1988.
 52. Wahlqvist ML, Kouris-Blazos A. *Diet related disorders—state of play*. Food and Nutrition Policy, Department of Community Services and Health, Canberra, Australia, 1991.
 53. US National Research Council. *Diet and health: implications for reducing chronic disease risk*. Washington DC: National Academy Press, 1989.
 54. Slater R, Kritchevsky D. *Human nutrition: a comprehensive treatise: cancer and nutrition*. New York: Plenum Press, 1991.
 55. Wahlqvist ML, Kouris-Blazos A. Olive-oil—nutrition and biological value. In: *World Encyclopedia of the Olive*. International Olive Oil Council, 1995.
 56. Frankel EN, Kanner J, German JB, Parks E, Kinsella KE. Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *Lancet* 1993;341:454-7.
 57. Byers TE. Diet and cancer: any progress in the interim? *Cancer* 1988;62:1713.
 58. Keys A, Menotti A, Karvonen MJ, et al. The diet and 15 year death rate in seven countries. *Am J Epidemiol* 1986;124:903.
 59. Horwarth CC. Dietary intake studies in elderly people. In: Bourne GH, ed *Impact of nutrition on health and disease*. *World Rev Nutr Diet* 1989;59:1-70.
 60. Wahlqvist ML, Kouris-Blazos A, Lukito W, Hsu-Hage B. Water soluble vitamin intakes in the elderly: cross-cultural findings in the IUNS study. In: Rosenberg IH, ed. *Nutritional assessment of elderly populations: measure and function*. New York: Raven Press, 1994;225-33.
 61. Roche AF. Anthropometric data. In: Wahlqvist ML, Davies L, Hsu-Hage BH-H, eds. *Cross-cultural study of food habits and health status in later life: description of elderly communities and lessons learnt*. *Asia Pacific J Clin Nutr* 1995 (in press).
 62. Chandra RK. The relation between immunology, nutrition and disease in elderly people. *Age Ageing* 1990;19(suppl):S25-31.
 63. Stadtman ER. Protein oxidation and ageing. *Science* 1992;257:1220-4.

Authors' addresses

A. Kouris-Blazos, M. L. Wahlqvist
Department of Medicine, Monash Medical Centre,
Monash University, Clayton Road, Clayton,
Melbourne, Victoria, Australia

A. Trichopoulou, E. Polychronopoulos
Department of Nutrition and Biochemistry,
Athens School of Public Health, Greece

D. Trichopoulos
Department of Epidemiology,
Harvard School of Public Health, USA

Received in revised form 29 September 1995